

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 13, 1994	3. REPORT TYPE AND DATES COVERED Annual Technical Report 9/1/93 - 8/31/94
4. TITLE AND SUBTITLE "Environmental Effects on High Energy Density Materials"			5. FUNDING NUMBERS 61103D 3484-XS
6. AUTHOR(S) Henry F. Schaefer III			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Georgia Univ Research Foundation Inc Boyd Grad Studies Res Cntr Athens, GA 30602			8. PERFORMING ORGANIZATION REPORT NUMBER AFOSR-TR- 94 0678
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/DENL Building 410, Bolling AFB DC 20332-6448 Dr Berman			10. SPONSORING / MONITORING AGENCY REPORT NUMBER F49620-93-J-0529
<div style="border: 2px solid black; padding: 5px; text-align: center;"> <p>DTIC SELECTED DEC 07 1994 F</p> </div>			
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED			12b. DISTRIBUTION CODE A
13. ABSTRACT (Maximum 200 words)  The <u>object</u> of this research is to characterize, using <u>ab initio</u> quantum mechanical methods, the stabilizing or destabilizing effects of hydrogen and/or oxygen matrices on proposed high energy density molecular (HEDM) systems. In addition, the unimolecular fragmentation reactions of large HEDM species such as N <sub>2</sub> O will be studied via density functional methods.			
14. SUBJECT TERMS			15. NUMBER OF PAGES
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT

19941201 100

*Technical Report*  
Air Force Office of Scientific Research  
Grant F49620-93-1-0529

"Environmental Effects on High Energy Density Materials"

*Principal Investigator: Henry F. Schaefer III*  
*Center for Computational Quantum Chemistry*  
*University of Georgia*

**I. Summary**

The development of efficient and safe conventional (i.e., nonnuclear) propellants and/or fuels is a goal of obvious technological significance. A desirable quality of such a propellant is clearly a high ratio of energy release to mass. The present hypothesis rests on a simple, but previously unrecognized, analogy between oxygen and sulfur. Preliminary studies showed that the oxygen ring systems are sufficiently promising to warrant the detailed, high-level theoretical research reported here.

Our idea begins with the observation that elemental sulfur exists as sulfur rings,  $S_n$ . The essence of our proposal is to make an analogy between sulfur rings and oxygen rings. Given the remarkable stability of sulfur rings, should it not be possible to prepare oxygen rings? Oxygen lies directly above sulfur in the Periodic Table, and the analogy is an appealing one.

Work completed during the past year (to September 1, 1994) concerns the prospective HEDM material  $C_2H_2Li_2$ . This molecule has been of sustained intellectual interest since 1976 when Paul Schleyer and John Pople suggested that the lowest triplet state of 1,1 -dilythioethylene might allow nearly free rotation about the supposed classical

<input checked="checked" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
Codes

Dist	Avail and/or Special
A-1	

C=C "double bond". In 1985 Manceron and Andrews intercepted dilithioethylene via matrix isolation infrared spectroscopy in the laboratory. Moreover, in 1987 Maercker, Graule, and Demuth used mercury precursors to characterize *cis* and *trans* 1,2-dilithioethylene as more conventional reaction products. So this species is by no means simply a figment of the theorist's imagination.

The work I describe was recently accepted for publication in the *Journal of the American Chemical Society*, having been carried out jointly with the Proctor and Gamble Company. The bulk of the research was carried out by my fourth year graduate student Evan Bolton, supported by the AASERT program.

The potential energy surface (PES) for the singlet 1,2-dilithioethene and acetylenic  $C_2H_2Li_2$  isomers was carefully surveyed using high level quantum mechanical methods. Three previously undiscovered minima (including, remarkably, the global minimum) were located: a planar monobridged *trans* 1,2 dilithioethene and two acetylenic structures. A total of seven minima and ten transition states for interconversion of minima were investigated, while seven transition states are located for the first time. Vibrational frequencies were evaluated for all structures through the coupled-cluster method including all single and double excitation with a double- $\zeta$  plus polarization basis set. A remarkable isomer, the  $C_s$  complex between lithioacetylene and LiH, is the global minimum on the  $C_2H_2Li_2$  PES. This structure was 34 kcal/mol more stable than the two lowest lying singlet 1,2-dilithioethene structures, a *trans* planar  $C_{2h}$  form with acute CCLi angles and a *cis* doubly bridged  $C_{2v}$  structure. The other singlet 1,2-dilithioethene minima, *cis* planar monobridged  $C_s$ , *cis* planar dibridged  $C_{2v}$ , and *trans* planar monobridged  $C_s$ , are 4.3, 8.4, and 19.4 kcal/mol higher lying, respectively. The carbon-lithium bonding is ionic in character in all these species.

## II. Publications (Since September 1, 1993)

1. E. E. Bolton, H. F. Schaefer, W. D. Laidig, and P. R. Schleyer, "Singlet  $C_2H_2Li_2$ : Acetylenic and 1, 2 - Dilithioethene Isomers. A Remarkably Congested Potential Energy Hypersurface for a Simple Organometallic System", *J. Amer. Chem. Soc.*, to appear in 1994.

### **III. List of Participating Professionals**

#### **A. Senior Research Personnel:**

Professor Henry F. Schaefer III

#### **B. Junior Research Personnel:**

Mr. Evan E. Bolton